

Exhibit B

```

<!--StartFragment-->RESULT 1
AAM52350
ID   AAM52350 standard; protein; 115 AA.
XX
AC   AAM52350;
XX
DT   25-JAN-2002 (first entry)
XX
DE   Extracellular domain of C-terminal lacking HM1.24 antigen.
XX
KW   HM1.24 antigen; extracellular domain; immunoassay; human.
XX
OS   Homo sapiens.
XX
PN   WO200177362-A1.
XX
PD   18-OCT-2001.
XX
PF   05-APR-2001; 2001WO-JP002964.
XX
PR   06-APR-2000; 2000JP-00105423.
XX
PA   (CHUS ) CHUGAI SEIYAKU KK.
XX
PI   Kinoshita Y, Ishikawa Y;
XX
DR   WPI; 2002-010919/01.
DR   N-PSDB; ABA01218.
XX
PT   Preparation of soluble HM1.24 antigen extracellular domain, useful for
PT   immunoassay of HM1.24 antigen.
XX
PS   Disclosure; Page 86-87; 96pp; Japanese.
XX
CC   The present invention relates to a method for preparing soluble HM1.24
CC   antigen extracellular domain. The method comprises culturing animal cells
CC   transformed by an expression vector carrying EF1alpha (undefined)
CC   promoter and HM1.24 antigen gene. The invention can be used for
CC   immunoassay of HM1.24 antigen. The method is more sensitive to
CC   conventional enzyme linked immunosorbant assay (ELISA) methods. The
CC   present sequence was used to illustrate the present invention
XX
SQ   Sequence 115 AA;

Query Match          100.0%; Score 574; DB 5; Length 115;
Best Local Similarity 100.0%; Pred. No. 4.7e-53;
Matches 115; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

Qy      1 NSEACRDGLRAVMECRNVTHLLQQELTEAQKGFQDVEAQAATCNHTVMALMASLDAEKAQ 60
        |||
Db      1 NSEACRDGLRAVMECRNVTHLLQQELTEAQKGFQDVEAQAATCNHTVMALMASLDAEKAQ 60

Qy      61 GQKKVEELEGEITTLNHKLQDASAEVERLRRENQVLSVRIADKKYYPSSQDSSSA 115
        |||
Db      61 GQKKVEELEGEITTLNHKLQDASAEVERLRRENQVLSVRIADKKYYPSSQDSSSA 115
<!--EndFragment-->

```